

APPROXIMATION PROPERTIES AP_s AND p-NUCLEAR OPERATORS (THE CASE WHERE $0 < s \leq 1$)

O. I. REINOV[†]

ABSTRACT. Among other things, it is shown that there exist Banach spaces Z and W such that Z^{**} and W have bases, and for every $p \in [1, 2)$ there is an operator $T : W \rightarrow Z$ that is not p -nuclear but T^{**} is p -nuclear.

We hold standard notation of the geometrical theory of operators in Banach spaces. The classical reference book on the theory of operator ideals is the A. Pietsch monograph [7]. Notations and terminology we use can be found, for instance, in [7], [8], [10], [11]. For our purposes, it is enough only to recall that if X, Y is a pair of Banach spaces and $p > 0$, then $N_p(X, Y)$ denotes the space of all p -nuclear operators from X to Y , and $X^* \widehat{\otimes}_p Y$ denotes the associated with it p -projective tensor product. And one more important reminder. If J is an operator ideal, then $J^{\text{reg}}(X, Y)$ denotes the space of all operators T from X into Y , for which $\pi_Y T \in J(X, Y^{**})$, where π_Y is the canonical isometric imbedding of the space Y into its second dual Y^{**} .

In this note we are interesting mainly in the following questions:

- 1) Under which conditions on Banach spaces X, Y the canonical mapping $X^* \widehat{\otimes}_p Y \rightarrow N_p(X, Y)$ is one-to-one?
- 2) Under which conditions on Banach spaces X, Y and on positive numbers s, r $N_s^{\text{reg}}(X, Y) \subset N_r(X, Y)$?

Both questions were investigated rather explicitly, e.g., in papers [10], [11]. The question 2) was considered also in [2] for $r = s = 1$ and in [8] for $r = s > 1$. For

[†] This work was done with partial support of the Ministry of the general and professional education of Russia (Grant 97-0-1.7-36) and FCP "Integracija", reg. No. 326.53.

[‡] AMS Subject Classification: 47B10. Hilbert–Schmidt operators, trace class operators, nuclear operators, p -summing operators, etc.

Key words and phrases: p -nuclear operators, bases, approximation properties, tensor products.

some other values of parameters p, r, s both questions were analyzed in the paper [3]. So, the well known A. Grothendieck's theorem on $2/3$ asserts that always one has $N_{2/3}^{\text{reg}} \subset N_1$ (somewhat below a proof of this assertion can be seen in the remark 2).

As more examples let us note also the following facts:

- a) if X, Y are arbitrary Banach spaces then the canonical mapping $X^* \widehat{\otimes}_{2/3} Y \rightarrow L(X, Y)$ is always one-to-one (see [3], [9], and also the corollary 1' below);
- b) if every finite-dimensional subspace E of a space Y is $C(\dim E)^\alpha$ -complemented in Y (where $0 < \alpha \leq 1/2$), then for $s, 1/s = 1 + \alpha$, and for every Banach space X the canonical mapping $X^* \widehat{\otimes}_s Y \rightarrow L(X, Y)$ is one-to-one (see [10]);
- c) if $0 < s \leq 1$ and $1/s = 1/2 + 1/r$, then $N_s^{\text{reg}}(X, Y) \subset N_r(X, Y)$ for any Banach spaces X, Y (see [3]);
- d) if $p \geq 1$ and $X^* \in \text{AP}$, then $N_p^{\text{reg}}(X, Y) \subset N_p(X, Y)$ for each Banach space Y (see [3] concerning the case $p = 1$; what about the case $p > 1$, here the proof is carried out by the same scheme that was in [3], so we are not going to repeat it).

Remark 1. A similar assertion as d) with assumption that $Y^{**} \in \text{AP}$ (instead of X^*) was formulated, without a proof, in [3] for $p = 1$ and in [5], with a proof based on [3], for $p > 1$. Below we will see that these last facts are not valid at least for the cases when $1 \leq p < 2$ (however, concerning the case $p = 1$ see the note [6]).

In [10], among others, it is shown that the assertions (a)–(c) can not be improved “in the scale of the spaces l_p .” Below we shall refine the assertions (a)–(c) for “the scale of Lorentz spaces”¹, and shall show that the new obtained results are unimprovable in this cases too.

We shall see also (and this is, perhaps, the main thing in the paper), that all the “negative” results of [8], [10], corresponding to the assertions (a)–(d), will take a place in the stronger versions. The first example of such a kind of the “sharpened” counterexamples may be found in the paper [6], where it was shown that there exists a Banach space Z with special properties: (i) Z^{**} is separa-

¹But not in the very full generality; below we shall be interesting in addition only in spaces $l_{s,\infty}$ for $s < 1$; but already in this case there are non entirely evident generalizations; see, e.g., theorem 4 below.

ble and has a basis (and Z^{***} does not possess the approximation property); (ii) $N_1^{\text{reg}}(Z^{**}, Z) \not\subset N_1(Z^{**}, Z)$. This is an example demonstrating that the second part of a Grothendieck's assertion [3] (ch.1, p. 86, proposition 15) is not valid. Below we will extend the mentioned counterexample to the case of the exponents of nuclearity p , bigger than one and less than two (but, certainly, differed from two; the corresponding counterexamples for $p > 2$ are the theme of a forthcoming paper); see the theorem 5 and its corollary 3.

Below we often shall use two following facts obtained in the paper [4], — the first one is exactly the theorem 1 in [4], and the second one is the corollary 1 from there (the existence of a basis in the space Z^{**} in the lemma 2 follows from the definition of the space Z in the proof of that corollary in [4]).

Lemma 1. *For each separable Banach space X there exist a separable Banach space E and a linear homomorphism $\psi : E^* \rightarrow X$ such that E^* has a basis and $E^{**} = \psi^*(X^*) \oplus \pi_E(E)$.*

Lemma 2. *For each separable Banach space Y there exist a separable Banach space Z and a linear homomorphism $\psi : Z^{**} \rightarrow Y$ such that Z^{**} has a basis (and, consequently, is separable), the kernel of the homomorphism ψ is equal to $\pi_Z(Z)$ and $\psi^*(Y^*)$ is complemented in Z^{***} .*

Corollary 1. *For each separable Banach space Y there exists a separable Banach space Z such that Z^{**} has a basis, $Z^{**}/\pi_Z(Z)$ is isomorphic to the space Y , $\pi_Z(Z)^\perp$ is complemented in Z^{***} and the following conditions are fulfilled:*

(a) *for every $p \in (0, 1]$ and every Banach space E the tensor product $E^* \widehat{\otimes}_p Y$ is isomorphic to the factor space $E^* \widehat{\otimes}_p (Z^{**}/\pi_Z(Z))$, which is, in turn, is isomorphic to*

$$N_p(E, Z^{**}) / \left[N_p^{\text{reg}}(E, \pi_Z(Z)) \cap N_1(E, \pi_Z(Z)) \right];$$

(b) *for every $p \in (0, 1]$ and every Banach space E the space $N_p(E, Y)$ is isomorphic to the space $N_p(E, Z^{**}/\pi_Z(Z))$, which is, in turn, is isomorphic to the factor space $N_p(E, Z^{**}) / N_p^{\text{reg}}(E, \pi_Z(Z))$.*

*In addition, all the isomorphisms in the assertions (a) and (b) are “canonical”, that is, are generated by the given homomorphism $Z^{**} \rightarrow Z^{**}/\pi_Z(Z) \rightarrow Y$ (details are in the proof).*

Proof. Let Z be the space from lemma 2. Denote, for the simplicity, $Z^{**}/\pi_Z(Z)$ by F .

Quotient map $\psi : Z^{**} \rightarrow Z^{**}/\pi_Z(Z) = F$ induces naturally the map $\Psi : N_p(E, Z^{**}) \rightarrow E^* \widehat{\otimes}_p F$, $\Psi(T) := \psi \circ T$. Since $p \leq 1$, it is clear that this map Ψ is a surjection. Let us see what is the kernel of this map.

Let $U \in \text{Ker } \Psi$. This means that for every $A \in L(F, E^{**})$ trace $A \circ \psi \circ U$ is equal to zero; or, that for each $B \in L(Z^{**}, E^{**})$ such that $B|_{\pi_Z(Z)} = 0$, trace $B \circ U$ is zero. It follows that $U(E) \subset \pi_Z(Z)$ (in other case one can find an operator $R \in L(Z^{**}, E^{**})$ such that $R|_{\pi_Z(Z)} = 0$ and trace $R \circ U \neq 0$).

So, $U \in N_p^{\text{reg}}(E, \pi_Z(Z))$. We have to refine this inclusion: let us prove that in fact $U \in N_1(E, \pi_Z(Z))$. Suppose that $U \notin N_1(E, \pi_Z(Z))$. Then one can find an operator $B \in L(Z^{**}, E^{**})$, such that trace $B \circ U = 1$, but trace $B \circ T = 0$ for every $T \in N_1(E, \pi_Z(Z))$. Since $\pi_Z(Z)$ has the Grothendieck approximation property, it follows now that $B|_{\pi_Z(Z)} = 0$, what is contradicted to the choice and properties of U . So, $U \in N_1(E, \pi_Z(Z))$. On the other hand, evidently, for every $V \in N_1(E, \pi_Z(Z))$ and $B \in L(Z^{**}, E^{**})$, where $B|_{\pi_Z(Z)} = 0$, we have: trace $B \circ V = 0$. Thus $\text{Ker } \Psi = N_p^{\text{reg}}(E, \pi_Z(Z)) \cap (N_1(E, \pi_Z(Z)))$. It follows from here the assertion (a).

Now consider the quotient map

$$\Psi_0 : N_p(E, Z^{**}) \xrightarrow{\Psi} E^* \widehat{\otimes}_p F \xrightarrow{j} N_p(E, F),$$

where j is the canonical quotient map. Its kernel $\text{Ker } \Psi_0$ consists of all the operator $U \in N_p(E, Z^{**})$ which are turned into the identical zero after the factorization $\psi : Z^{**} \rightarrow F$, that is of those U , for which $U(E) \subset \text{Ker } \psi = \pi_Z(Z)$. This means that $\text{Ker } \Psi_0 = N_p^{\text{reg}}(E, \pi_Z(Z))$. ■

Let us note a curious “auxiliary” effect of our arguments (this is, certainly, something like “to break a butterfly on the wheel”):

Corollary 1' (A. Grothendieck [3]). *If $p \in (0, 2/3]$ then $G^* \widehat{\otimes}_p Y = N_p(G, Y)$ for every Banach spaces G and Y (that is there is no factorization under the canonical quotient map $G^* \widehat{\otimes}_p Y \rightarrow N_p(G, Y)$).*

Proof. Clearly, we can consider only separable Banach spaces². If $p \leq 2/3$ then every operator $U \in N_p^{\text{reg}}(G, \pi_Z(Z))$, where Z is from lemma 2 (or, that is the

²For each separable $H \subset G^*$ there exists a separable subspace $G_0 \subset G$ such that for the identity imbedding $j : G_0 \rightarrow G$ the operator $j^*|_H : G_0^* \rightarrow G_0^*$ is an isometric imbedding.

same, from corollary 1), factors through a nuclear diagonal operator (see below remark 2) and, therefore, is nuclear itself. Thus $N_p^{\text{reg}}(G, \pi_Z(Z)) \cap (N_1(G, \pi_Z(Z))) = N_p^{\text{reg}}(G, \pi_Z(Z))$ and the corollary 1 is applied. ■

Recall that a Banach space X has the approximation property of the order p , $p \in (0, +\infty]$ (briefly, $X \in \text{AP}_p$), if for every Banach space Z the canonical mapping $Z^* \widehat{\otimes}_p X \rightarrow L(Z, X)$ is one-to-one (see, e.g., [8], [10]). It is convenient to introduce some more additional notation and definition connected with the approximation properties. Namely, for a pair of the spaces X, Y and a number $s \in (0, 1)$ we denote by $Y^* \widehat{\otimes}_{s, \infty} X$ the linear space consisting of tensors $z \in Y^* \widehat{\otimes}_1 X$ (the projective tensor product), admitted representations of the kind

$$z = \sum_{k=1}^{\infty} \lambda_k y'_k \otimes x_k \quad \text{where} \quad \lambda_k \searrow, \lambda_k^s = o(1/k), \|y'_k\| \|x_k\| = 1.$$

Definition 1. The Banach space X has the property $\text{AP}_{s, \infty}$, where $s \in (0, 1)$, if for each Banach space Z the canonical mapping $Z^* \widehat{\otimes}_{s, \infty} X \rightarrow L(Z, X)$ is one-to-one (in other words, $Z^* \widehat{\otimes}_{s, \infty} X = N_{s, \infty}(Z, X)$; it must be clear what the operator space on the right is).

Note that a space X has the property $\text{AP}_{s, \infty}$ iff the canonical mapping $X^* \widehat{\otimes}_{s, \infty} X \rightarrow L(X, X)$ is one-to-one; in this case $X^* \widehat{\otimes}_{s, \infty} X = N_{s, \infty}(X, X)$.

Theorem 1. 1) Let $\alpha \in (0, 1/2]$, $C > 0$, and X be a Banach space. If every finite-dimensional subspace E of the space X is $C(\dim E^\alpha)$ -complemented in X then $X \in \text{AP}_{s, \infty}$, where $1/s = 1 + \alpha$. In particular (take $\alpha = 1/2$), every Banach space has the property $\text{AP}_{2/3, \infty}$, and if X is a subspace or a factor space of some space $L^p(\mu)$, where $p \in (1, +\infty)$, $p \neq 2$, then $X \in \text{AP}_{s, \infty}$ ($1/s = 1 + |1/p - 1/2|$).

2) For each $s \in [2/3, 1)$ there exists a separable reflexive space Y such that $Y \in \text{AP}_{s, \infty}$, but $Y \notin \text{AP}_r$ for any $r \in (s, 1]$.

The *Proof* of the part 2) of the theorem is completely analogous to the proof of the corresponding fact from [10] (namely, the assertions 1) and 2) of the theorem 5.4 in [10]); the proof of this theorem 5.4 is literally carried over to our case, but instead of the lemma 5.1 from [10] one has to use such a strengthening of the part 1 of our theorem 1:

Proposition 1. *Let $\alpha \in (0, 1/2]$, $C > 0$, and X be a Banach space. Suppose that for every finite-dimensional subspace E of the space X one can find a finite-dimensional subspace F , $E \subset F \subset X$, such that F is $C(\dim E)^\alpha$ -complemented in X . Then $X \in \text{AP}_{s,\infty}$, where $1/s = 1 + \alpha$.*

Again, it is enough to refer to the paper [10]. Our Proposition 1 is proved implicitly in [10], Lemma 5.1: if we look carefully at the proof of the lemma in [10], then it is not hard to see that the main thing used in that proof is the belonging of the tensor z to the projective tensor product (in this paper all the tensors have this property) and the estimation $\lambda_j^s = o(j^{-1})$ for the coefficients of z from the proof of the mentioned Lemma 5.1. So, we will not repeat all those arguments from [10] which were used in the proof of this lemma, and refer the reader to [10].

Theorem 2. 1) *Let $s \in (0, 1)$. If $X^* \in \text{AP}_{s,\infty}$ or $Y^{***} \in \text{AP}_{s,\infty}$, then $N_{s,\infty}^{\text{reg}}(X, Y) \subset N_1(X, Y)$. In particular, for every Banach spaces X and Y we have: $N_{2/3,\infty}^{\text{reg}}(X, Y) \subset N_1(X, Y)$.*

2) *There exists a Banach space Z with the following properties:*

- (i) *Z^{**} has a basis (therefore, $Z^{**} \in \text{AP}$);*
- (ii) *for each $s \in (2/3, 1]$ $N_s^{\text{reg}}(Z^{**}, Z) \not\subset N_1(Z^{**}, Z)$.*

Thus, the approximation conditions imposed on X and Y in 1) are essential.

Proof. 1) Suppose there exists such an operator $T \in L(X, Y)$ that $T \notin N_1(X, Y)$ but $\pi_Y T \in N_{s,\infty}(X, Y^{**})$. Since either X^* or Y^{**} has the property $\text{AP}_{s,\infty}$, we have:

$$N_{s,\infty}(X, Y^{**}) = X^* \widehat{\otimes}_{s,\infty} Y^{**}.$$

Hence the operator $\pi_Y T$ can be identified with a tensor element $t \in X^* \widehat{\otimes}_{s,\infty} Y^{**} \subset X^* \widehat{\otimes}_1 Y^{**}$; in addition, by the choice of T , $t \notin X^* \widehat{\otimes}_1 Y$ (as usually, $X^* \widehat{\otimes}_1 Y$ is considered as a subspace of the space $X^* \widehat{\otimes}_1 Y^{**}$). Therefore, there exists such an operator $U \in L(Y^{**}, X^{**}) = (X^* \widehat{\otimes}_1 Y^{**})^*$ that $\text{trace } U \circ t = \text{trace } (t^* \circ (U^*|_{X^*})) = 1$ and $\text{trace } U \circ \pi_Y \circ z = 0$ for each $z \in X^* \widehat{\otimes}_1 Y$. It follows now that $U\pi_Y = 0$ and $\pi_Y^* U^*|_{X^*} = 0$. Indeed, if $x' \in X^*$ and $y \in Y$ then

$$\langle U\pi_Y y, x' \rangle = \langle y, \pi_Y^* U^*|_{X^*} x' \rangle = \text{trace } U \circ (x' \otimes \pi_Y(y)) = 0.$$

Evidently the tensor element $U \circ t$ induces the operator $U\pi_Y T$, which is identically equal to zero.

If $X^* \in \text{AP}_{s,\infty}$ then $X^* \widehat{\otimes}_{s,\infty} Y^{**} = N_{s,\infty}(X, Y^{**})$, hence this tensor element is zero what is contradicted to the equality $\text{trace } U \circ t = 1$.

Let now $Y^{***} \in \text{AP}_{s,\infty}$. In this case the operator

$$V := (U^*|_{X^*}) \circ T^* \circ \pi_Y^* : Y^{***} \rightarrow Y^* \rightarrow X^* \rightarrow Y^{***}$$

uniquely defines a tensor element $t_0 \in Y^{***} \widehat{\otimes}_{s,\infty} Y^{***}$. Take any representation $t = \sum x'_n \otimes y''_n$ for t as an element of the space $X^* \widehat{\otimes}_{s,\infty} Y^{**}$. Denoting for the brevity the operator $U^*|_{X^*}$ by U_* , we get:

$$\begin{aligned} Vy''' &= U_* (T^* \pi_Y^* y''') = U_* ((T^* \pi_Y^* \pi_{Y^*}) \pi_Y^* y''') = U_* ((\pi_Y T)^* \pi_{Y^*}) \pi_Y^* y''' = \\ &= U_* \left(\left(\sum y''_n \otimes x'_n \right) \pi_{Y^*} \pi_Y^* y''' \right) = U_* \left(\sum \langle y''_n, \pi_Y^* y''' \rangle x'_n \right) = \\ &= \sum \langle \pi_Y^{**} y''_n, y''' \rangle U_* x'_n. \end{aligned}$$

So, the operator V (or the element t_0) has in the space $Y^{***} \widehat{\otimes}_{s,\infty} Y^{***}$ the representation

$$V = \sum \pi_Y^{**}(y''_n) \otimes U_*(x'_n).$$

Therefore

$$\text{trace } t_0 = \text{trace } V = \sum \langle \pi_Y^{**}(y''_n), U_*(x'_n) \rangle = \sum \langle y''_n, \pi_Y^* U_* x'_n \rangle = \sum 0 = 0.$$

On the other hand,

$$Vy''' = U_* (\pi_Y T)^* y''' = U_* \circ t^*(y''') = U_* \left(\sum \langle y''_n, y''' \rangle x'_n \right) = \sum \langle y''_n, y''' \rangle U_* x'_n,$$

whence $V = \sum y''_n \otimes U_*(x'_n)$. Therefore

$$\text{trace } t_0 = \text{trace } V = \sum \langle y''_n, U_* x'_n \rangle = \sum \langle U y''_n, x'_n \rangle = \text{trace } U \circ t = 1.$$

The obtained contradiction completes the proof of the first part of 1). The case $s = 3/2$ is reduced to the just proved assertion by using the first part of Theorem 1.

2) It is well known that there exist a separable reflexive Banach space X and a nonzero tensor element $z \in X^* \widehat{\otimes}_1 X$ such that $z \in X^* \widehat{\otimes}_s X$ for all $s > 2/3$, but the associated operator $\tilde{z}$ is identically zero (see, e.g., [1], [7]). Thus for each $s \in (0, 1]$ $X^* \widehat{\otimes}_s X \neq N_s(X, X)$. By Corollary 1 there are a separable Z and a homomorphism

$\varphi : Z^{**} \rightarrow X$ such that Z^{**} has a basis, $\text{Ker } \varphi = \pi_Z(Z)$, $\varphi^*(X^*)$ is complemented in Z^{***} and $N_s^{\text{reg}}(X, Z) \not\subset N_1(X, Z)$ for all $s > 2/3$. Since $\varphi^*(X^*)$ is complemented in Z^{***} , then $N_s^{\text{reg}}(Z^{**}, Z) \not\subset Z^{***} \widehat{\otimes}_1 Z = N_1(Z^{**}, Z)$ for each $s > 2/3$. ■

Remark 2. It is rather easy to show that $N_{2/3}^{\text{reg}} \subset N_1$. Indeed, if an operator T acts from a space E to a space Z and is $2/3$ -nuclear as an operator from E into Z^{**} then, by the very definition of the $2/3$ -nuclear operator, “splitting” the coefficients of its tensor-series expansion into pairs of appropriately chosen factors, we can expand the operator T itself into a product as follows:

$$T : E \xrightarrow{A} c_0 \xrightarrow{\Delta_1} l^1 \xrightarrow{j} l^2 \xrightarrow{\Delta_2} l^1 \xrightarrow{B} Z^{**}$$

where Δ_j are diagonal operators (thus, Δ_1 is nuclear) and other operators are continuous. Taking an orthogonal projection in l^2 onto the closure of the image of the operator $j \Delta_1 A$, and considering B only on this subspace, we get the same operator $T : E \rightarrow Z$; this simple trick shows that T is nuclear from E into Z . At this moment I have no idea how to obtain the inclusion $N_{2/3, \infty}^{\text{reg}} \subset N_1$ directly by an analogous way (maybe, one have to get a factorization through l_2 not for whole operator but for its finite dimensional parts by using the fact that every n -dimensional subspace of a Banach space is $n^{1/2}$ -complemented).

Corollary 2. *Let X be such a reflexive Banach space that each its n -dimensional subspace Cn^α -complemented in X (here $C > 0, \alpha \in (0, 1/2)$ are some constances, $n = 1, 2, \dots$). Then for every Banach space Y*

$$N_{s, \infty}^{\text{reg}}(X, Y) \subset N_1(X, Y).$$

where $1/s = 1 + \alpha$.

In particular, if $X \subset L^p(\mu), 1 < p < +\infty, p \neq 2, \quad 1/s = 1 + |1/2 - 1/p|$, then

$$N_{s, \infty}^{\text{reg}}(X, Y) \subset N_1(X, Y) \quad \text{for every } Y.$$

To prove the corollary it is enough to apply Theorem 1,1) and Theorem 2,1) and to use the fact that any n -dimensional subspace of $L^p(\mu)$ $Cn^{|1/2 - 1/p|}$ -complemented in $L^p(\mu)$.

Remark 3. In Theorem 2,1) and Corollary 2 we obtained even the following inclusion: $N_{s, \infty}^{\text{reg}}(X, Y) \subset X^* \widehat{\otimes}_1 Y$. One has to understand this in the following manner:

every operator $T \in N_{s,\infty}^{\text{reg}}(X, Y)$ is uniquely generated by a tensor element from the space $X^* \widehat{\otimes}_{s,\infty} Y^{**} \subset X^* \widehat{\otimes}_1 Y^{**}$, which (the tensor element), in turn, belongs to the space $X^* \widehat{\otimes}_1 Y$, a subspace of $X^* \widehat{\otimes}_1 Y^{**}$; so if

$$\begin{aligned} \sum x'_n \otimes y''_n & \text{ is a representation of } T \text{ in the space } X^* \widehat{\otimes}_{s,\infty} Y^{**}, \\ \sum \tilde{x}'_n \otimes \tilde{y}''_n & - \text{ in the space } X^* \widehat{\otimes}_1 Y^{**} \text{ and} \\ \sum \tilde{\tilde{x}}'_n \otimes \tilde{\tilde{y}}_n & - \text{ in the space } X^* \widehat{\otimes}_1 Y, \end{aligned}$$

then for each $U \in L(Y^{**}, X^{**})$

$$\text{trace } U \circ T = \sum \langle x'_n, U y''_n \rangle = \sum \langle \tilde{x}'_n, U \tilde{y}''_n \rangle = \sum \langle \tilde{\tilde{x}}'_n, U \tilde{\tilde{y}}_n \rangle$$

(let us note that the trace is well defined since in Theorem 2,1) and Corollary 2 the natural map $N_{s,\infty}^{\text{reg}}(X, Y^{**}) \rightarrow L(X, Y^{**})$ is one-to-one).

By the same method as in the proof of Theorem 2,1) we get

Theorem 2'. *Let $s \in (0, 1]$. If $X^* \in \text{AP}_s$, or $Y^{***} \in \text{AP}_s$ then $N_s^{\text{reg}}(X, Y) \subset N_1(X, Y)$.*

Looking at the proof of the part 2) of Theorem 2 and of Corollary 2 it is not hard to see that we have the following stronger result:

Proposition 2. *There exist a separable Banach space Z and an operator $A \in L(Z^{**}, Z)$ so that Z^{**} has a basis, $\pi A \in N_s(Z^{**}, Z^{**})$ for all $s > 2/3$ but $A \notin N_1(Z^{**}, Z)$.*

Theorem 3. *For each $s \in [2/3, 1)$ there exists a separable Banach space Z so that*

- (i) Z^{**} has a basis;
- (ii) all the spaces $Z, Z^*, Z^{**}, \dots$ have the property AP_s ;
- (iii) Z^{***} does not have the property AP_r if $r \in (s, 1]$;
- (iv) for every Banach space X

$$N_{s,\infty}^{\text{reg}}(X, Z) \subset N_1(X, Z) \quad \text{and} \quad N_{s,\infty}^{\text{reg}}(Z^{**}, X) \subset N_1(Z^{**}, X);$$

- (v) for each $r \in (s, 1]$ $N_r^{\text{reg}}(Z^{**}, Z) \not\subset N_1(Z^{**}, Z)$. Moreover, there exists an operator $U : Z^{**} \rightarrow Z$ such that $U \in N_r^{\text{reg}}(Z^{**}, Z)$ for all $r \in (s, 1]$, but $U \notin N_1(Z^{**}, Z)$.

The method of the proof of Theorem 3 is essentially the same as one for the part 2) of Theorem 2 by using Theorem 1,2), Corollary 1 and Theorem 2,1). We omit the proof leaving it to the reader as an not difficult exercise.

Let us look now at the situation when there are no approximation restrictions on Banach spaces under consideration (as, for instance, in the case $s = 2/3$ in Theorem 2,1) and pose the question of what assertion can be obtained instead of one in the part 1) of Theorem 2.

The following fact we shall use is proved essentially in [10] (Theorem 2,1,A). It is convenient to denote by φ_{sp}^∞ the canonical map $X^* \widehat{\otimes}_{s,\infty} Y \rightarrow X^* \widehat{\otimes}_p Y$ and by j_p the canonical map $X^* \widehat{\otimes}_p Y \rightarrow L(X, Y)$.

Proposition 3. *If $s \leq 1$ and $p \geq 2s/(2-s)$ then the mapping*

$$i_p : \varphi_{sp}^\infty (X^* \widehat{\otimes}_{s,\infty} Y) \rightarrow L(X, Y)$$

is one-to-one.

One can find a proof of this proposition in [10] (see there the proof of the part A of Theorem 2,1).

The following fact is a stronger version of the theorem 3.1),A) from [10] where it was shown that if s and p are as in Proposition 3 then $N_s^{\text{reg}}(X, Y) \subset N_p(X, Y)$. However, the proof will be not so elementary as in [10].

Theorem 4. *If $2/3 \leq s < 1$ and $p = 2s/(2-s)$ then*

$$N_{s,\infty}^{\text{reg}}(X, Y) \subset N_p(X, Y)$$

for any Banach spaces X and Y .

Proof. Let $T \in N_{s,\infty}^{\text{reg}}$ and $T \neq 0$. Fix a tensor element $t = \sum \bar{x}'_n \otimes \bar{y}''_n \in X^* \widehat{\otimes}_{s,\infty} Y^{**}$, which generates the operator T (i.e., $\pi_Y T = \tilde{t}$). It is enough to prove that $t \in X^* \widehat{\otimes}_p Y$ (i.e, that the image $\varphi_{sp}^\infty(t) \in X^* \widehat{\otimes}_p Y^{**}$ lies in fact in the subspace $X^* \widehat{\otimes}_p Y$).

Suppose that $t \notin X^* \widehat{\otimes}_p Y$. In this case one can find an operator $U \in \Pi_{p'}(Y^{**}, X^{**})$, for which $\text{trace } U \circ t = 0$ and, on the other hand, $\text{trace } U \circ z = 1$ for all $z \in X^* \widehat{\otimes}_p Y$. Further on it will be convenient to consider several diagrams (with some comments):

$$X \xrightarrow{T} Y \xrightarrow{\pi_Y} Y^{**} \xrightarrow{B} Z_{p'} \xrightarrow{S} X^{**} \xrightarrow{T^{**}} \pi_Y(Y)$$

($U = SB$; $\tilde{t} = \pi_Y T$; $B\pi_Y = jB_0$ and $A := \pi^{-1}T^{**}S$ (these operators will appear later); $U\pi_Y = 0$ and $Sj = 0$);

$$Y \xrightarrow{B_0} Y_{p'} \xrightarrow{j} Z_{p'} \xrightarrow{S} X^{**} \xrightarrow{T^{**}} \pi_Y(Y) \xrightarrow{\pi_Y^{-1}} Y \xrightarrow{B_0} Y_{p'}.$$

Here $Z_{p'} \subset L_{p'}(\mu)$ (for some finite measure μ); $B \in \Pi_{p'}(Y^{**}, Z_{p'})$ and $S \in L(Z_{p'}, X^{**})$ are such that $U = SB$; $Y_{p'} = B\pi_Y(Y)$; B_0 is the operator induced by B ; j is the identity imbedding. Note that $T^{**}(X^{**}) \subset \pi_Y(Y)$ (by the compactness of T). Besides, $U\pi_Y = 0$ by the assumption; hence, $Sj = 0$. Denote the operator $\pi_Y^{-1}T^{**}S$ by A .

Since $Z_{p'} \subset L_{p'}(\mu)$ and $\pi_Y A \in N_{s,\infty}$, then, by Theorem 1,1), the operator $\pi_Y A$ is uniquely defined by a tensor element from $Z_{p'}^* \widehat{\otimes}_{s,\infty} Y^{**}$, which will be denoted again by $\pi_Y A$. It is clear that

$$\pi_Y A = T^{**}S = \sum S^* (\pi_{X^*}(\bar{x}'_n)) \otimes \bar{y}''_n.$$

By Corollary 2 and Remark 3, $A \in Z_{p'}^* \widehat{\otimes}_1 Y$, and also

$$\begin{aligned} \text{trace } B(\pi_Y A) &= \text{trace } (B\pi_Y) \circ A = \sum \langle S^* \pi_{X^*} \bar{x}'_n, B\bar{y}''_n \rangle = \\ &= \sum \langle \bar{x}'_n, U\bar{y}''_n \rangle = \text{trace } U \circ t = 1. \end{aligned}$$

In particular, it follows from here that $\pi_Y A \neq 0$ (and $A \neq 0$).

Take a representation of A of the kind $\sum z'_n \otimes y_n$ as an element of the space $Z_{p'}^* \otimes_1 Y$; this is also a representation of the tensor element $\pi_Y A \in Z_{p'}^* \widehat{\otimes}_{s,\infty} Y^{**}$ (but in the space $\pi_Y A \in Z_{p'}^* \widehat{\otimes}_1 Y^{**}$). We have:

$$\begin{aligned} 1 &= \text{trace } (B\pi_Y) \circ A = \text{trace } (jB_0) \circ A = \text{trace } (jB_0) \circ \left(\sum z'_n \otimes y_n \right) = \\ &= \sum \langle z'_n, jB_0 y_n \rangle = \sum \langle z'_n, jB_0^{**} \pi_Y y_n \rangle = \text{trace } (jB_0^{**}) \circ \left(\sum z'_n \otimes \pi_Y y_n \right) = \\ &= \text{trace } (jB_0^{**}) \circ (\pi_Y A) = \text{trace } (jB_0^{**}) \circ \left(\sum S^* (\pi_{X^*}(\bar{x}'_n)) \otimes \bar{y}''_n \right) = \\ &= \sum \langle S^* \pi_{X^*} \bar{x}'_n, jB_0^{**} \bar{y}''_n \rangle = \text{trace } B_0^{**} \circ \left(\sum ((Sj)^* \pi_{X^*} \bar{x}'_n) \otimes \bar{y}''_n \right). \end{aligned}$$

It follows from here that the tensor element $\alpha := \sum ((Sj)^* \pi_{X^*} \bar{x}'_n) \otimes \bar{y}''_n$, belonging to the space $Y_{p'}^* \widehat{\otimes}_{s,\infty} Y^{**}$, is not equal to zero. By Theorem 1,1), the associated operator $\tilde{\alpha}$ is not zero too. It is remain to note that $\tilde{\alpha} = T^{**}Sj = 0$. $\blacksquare$

Remark 4. The assertion A' of Theorem 3.1 from [10] shows that Theorem 4 is sharp. An analogous remark can be made about Proposition 3. However the following result is much stronger than the part A' of the theorem 3.1 in [10] where it is obtained that there are two Banach spaces X and Y such that, in particular, $N_s^{\text{reg}}(X, Y) \not\subset N_p(X, Y)$ and $Y \in \text{AP}$. Having in mind the result of A. Grothendieck (which we said about before the definition 1 and which is now clear to be partially wrong – see [6]) I wrote in [10] that the space Y "certainly does not have the BAP". Now we shall see that this last phrase was not reasonable.

Theorem 5. *Let $r \in [2/3, 1)$ and $1/r = 1/2 + 1/p$. There exist two separable Banach spaces Z and W such that*

- (i) W and Z^{**} have bases;
- (ii) all the duals of the space W have the property $\text{AP}_{r, \infty}$;
- (iii) W^* does not have AP_s for any $s \in (r, 1]$;
- (iv) for each Banach space E $N_{r, \infty}^{\text{reg}}(W, E) \subset N_1(W, E)$;
- (v) $N_s^{\text{reg}}(W, Z) \not\subset N_p(W, Z)$ for any $s \in (r, 1]$.

Proof. It is enough to prove that for every $s, r < s < 1$, there are Banach spaces $Z = Z_s$ and $W = W_s$ with the properties mentioned in the statement of the theorem; moreover, if s is fixed, it is enough to get in (iii) the weaker condition $W^* \notin \text{AP}_{s, \infty}$ (since s is running over the interval $(r, 1)$, and the property AP_1 is the strongest one among all of the approximation properties). If we do this then for $s_n \searrow r$ the spaces $(\sum Z_{s_n})_{l_2}$ and $(\sum W_{s_n})_{l_2}$ will be completely satisfied all the conditions of the theorem.

Thus, let $s \in (r, 1)$, so $p < 2s/(2 - s)$. By Theorem 2.1, A'), 1) in [10], there exist separable reflexive spaces X and Y , a tensor element $z \in X^* \widehat{\otimes}_s Y$ and an operator $U \in \mathcal{QN}_{p'}(Y, X)$ (quasi- p' -nuclear) such that $\text{trace } U \circ z = 1$ and the associated operator $\tilde{z} = 0$; moreover, the space X can be taken as a subspace of a space $L_{p'}$ (the last fact is contained explicitly in the proof of the mentioned theorem from [10]).

Let Z be such a separable space that Z^{**} has a basis and there is a homomorphism φ from Z^{**} onto Y with the kernel $\pi_Z(Z)$ (see Lemma 2). Lift up the (nuclear) tensor element z , lying in $X^* \widehat{\otimes}_s Y$, to get an element $\alpha \in X^* \widehat{\otimes}_s Z^{**}$, so that $\varphi \circ \alpha = z$, and set $V := U \circ \varphi$. Since $\text{trace } V \circ \alpha = \text{trace } U \circ z = 1$ and Z^{**} possesses the

approximation property (the property AP_1), then $\tilde{\alpha} = \alpha \neq 0$. Besides, the operator $\widehat{\varphi \circ \alpha}$, associated with the tensor $\varphi \circ \alpha$, is equal to zero. Therefore $\alpha(X) \subset \text{Ker } \varphi = \pi_Z(Z)$, that is the operator α acts from X to Z .

If $\alpha \in X^* \widehat{\otimes}_p Z$, then for any its p -nuclear representation $\alpha = \sum x'_n \otimes z_n$ we have: $\text{trace } V \circ \alpha = \sum \langle x'_n, Vz_n \rangle = \sum 0 = 0$; hence $\alpha \notin X^* \widehat{\otimes}_p Z$ (we used the fact that $V \in \Pi_{p'}(Z, X)$). Thus $N_s^{\text{reg}}(X, Z) \not\subset N_p(X, Z)$.

Let us note that X (and X^*) has the property $AP_{r,\infty}$ (Theorem 1,1) — remember where the space X lies), but does not have the property $AP_{s,\infty}$ (Theorem 2,1)).

Now, consider such a separable Banach space E that E^* has a basis and there is a homomorphism ψ from E^* onto X with the property that $E^{**} = \psi^*(X^*) \oplus \pi_E(E)$ (Lemma 1). Set $W = E^*$. Since $E \in AP_1$ and $X^* \in AP_{r,\infty}$, all the conjugate spaces $W^*, W^{**}, \dots$ possess the property $AP_{r,\infty}$; by Theorem 2,1), the assertion (iv) is fulfilled, and by the construction the same is true for (v). Besides, $W^* \notin AP_{s,\infty}$. ■

It follows from (i) and (v) of the previous theorem

Corollary 3. *There exists a pair of separable Banach spaces Z, W with the following properties. The spaces Z^{**} and W have bases (therefore, have the approximation property), and for every $p, 1 \leq p < 2$, one can find non- p -nuclear operator from W to Z with p -nuclear second adjoint.*

Remark 5. Such a phenomenon is impossible if either W^* or Z^{***} has the approximation property. This was mentioned above in passing (a particular case was formulated in Theorem 2'). As to the case $p > 2$, it is the theme of the forthcoming paper of the author.

REFERENCES

1. A. M. Davie, *The approximation problem for Banach spaces*, Bull. London Math. Soc. **5** (1973), 261-266.
2. T. Figiel, W. B. Johnson, *The approximation property does not imply the bounded approximation property*, Proc. Amer. Math. Soc. **41** (1973), 197-200.
3. A. Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, Mem. Amer. Math. Soc. **16** (1955), 196-140.

4. J. Lindenstrauss, *On James' paper "Separable Conjugate Spaces"*, Israel J. Math. **9** (1971), 279-284.
5. B. M. Makarov, V. G. Samarskii, *Weak sequential completeness and close properties of some spaces of operators*, in book "Theory of operators and theory of functions". Leningrad University (1983), 122-144, (in Russian).
6. E. Oja, O. I. Reinov, *Un contre-exemple à une affirmation de A. Grothendieck*, C. R. Acad. Sc. Paris. — Serie I **305** (1987), 121-122.
7. A. Pietsch, *Operator ideals*, North-Holland (1980).
8. O. I. Reinov, *Approximation properties of order p and the existence of non- p -nuclear operators with p -nuclear second adjoints*, Math. Nachr. **109** (1982), 125-134.
9. O. I. Reinov, *A simple proof of two theorems of A. Grothendieck*, Vestnik LGU **7** (1983), 115-116, (in Russian).
10. O. I. Reinov, *Disappearing tensor elements in the scale of p -nuclear operators*, in book "Theory of operators and theory of functions". Leningrad University (1983), 145-165, (in Russian).
11. O. I. Reinov, *Sur les operateurs p -nucléaires entre espaces de Banach avec bases*, C. R. Acad. Sc. Paris. — Serie I **316** (1993), 905-907.

OLEG I. REINOV
 DEPARTMENT OF MATHEMATICS
 ST PETERSBURG UNIVERSITY
 ST PETERHOF, BIBLIOTEC PL 2
 198904 ST PETERSBURG, RUSSIA
E-mail address: orein@orein.usr.pu.ru